



## ALDH5A1 gene

aldehyde dehydrogenase 5 family member A1

### Normal Function

The *ALDH5A1* gene provides instructions for producing the succinic semialdehyde dehydrogenase enzyme. This enzyme is found in the energy-producing centers of cells (mitochondria). Succinic semialdehyde dehydrogenase is involved in the breakdown of a chemical that transmits signals in the brain (neurotransmitter) called gamma-amino butyric acid (GABA). The primary role of GABA is to prevent the brain from being overloaded with too many signals. Once GABA molecules have been released from nerve cells, they are broken down by succinic semialdehyde dehydrogenase and other enzymes.

### Health Conditions Related to Genetic Changes


#### succinic semialdehyde dehydrogenase deficiency

At least 35 mutations in the *ALDH5A1* gene have been found to cause succinic semialdehyde dehydrogenase deficiency. Most of these mutations change one protein building block (amino acid) in the succinic semialdehyde dehydrogenase enzyme. Mutations in the *ALDH5A1* gene lead to the production of an enzyme with little or no activity. A lack of functional succinic semialdehyde dehydrogenase disrupts the conversion of succinic semialdehyde to succinic acid. Instead, succinic semialdehyde is converted back into GABA or to a related molecule, gamma-hydroxybutyrate (GHB). It is unclear how increases in GHB and GABA cause developmental delay, seizures, and other features of succinic semialdehyde dehydrogenase deficiency.

## Chromosomal Location

Cytogenetic Location: 6p22.3, which is the short (p) arm of chromosome 6 at position 22.3

Molecular Location: base pairs 24,494,969 to 24,537,207 on chromosome 6 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

## Other Names for This Gene

- aldehyde dehydrogenase 5 family, member A1
- aldehyde dehydrogenase 5 family, member A1 (succinate-semialdehyde dehydrogenase)
- aldehyde dehydrogenase 5A1
- mitochondrial succinate semialdehyde dehydrogenase
- NAD(+)-dependent succinic semialdehyde dehydrogenase
- SSADH
- SSDH
- SSDH\_HUMAN

## Additional Information & Resources

### Educational Resources

- Basic Neurochemistry (sixth edition, 1999): GABA shunt reactions are responsible for the synthesis, conservation and metabolism of GABA  
<https://www.ncbi.nlm.nih.gov/books/NBK27979/?rendertype=figure&id=A1180>
- Basic Neurochemistry (sixth edition, 1999): GABA Synthesis, Uptake and Release  
<https://www.ncbi.nlm.nih.gov/books/NBK27979/>

### GeneReviews

- Succinic Semialdehyde Dehydrogenase Deficiency  
<https://www.ncbi.nlm.nih.gov/books/NBK1195>

### Scientific Articles on PubMed

- PubMed  
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28ALDH5A1%5BTIAB%5D%29+OR+%28%28SSDH%5BTIAB%5D%29+OR+%28SSADH%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+2880+days%22%5Bdp%5D>

### OMIM

- ALDEHYDE DEHYDROGENASE 5 FAMILY, MEMBER A1  
<http://omim.org/entry/610045>

### Research Resources

- ClinVar  
<https://www.ncbi.nlm.nih.gov/clinvar?term=ALDH5A1%5Bgene%5D>
- HGNC Gene Family: Aldehyde dehydrogenases  
<http://www.genenames.org/cgi-bin/genefamilies/set/398>
- HGNC Gene Symbol Report  
[http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?q=data/hgnc\\_data.php&hgnc\\_id=408](http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=408)
- NCBI Gene  
<https://www.ncbi.nlm.nih.gov/gene/7915>
- UniProt  
<http://www.uniprot.org/uniprot/P51649>

### **Sources for This Summary**

- OMIM: ALDEHYDE DEHYDROGENASE 5 FAMILY, MEMBER A1  
<http://omim.org/entry/610045>
- Akaboshi S, Hogema BM, Novelletto A, Malaspina P, Salomons GS, Maropoulos GD, Jakobs C, Grompe M, Gibson KM. Mutational spectrum of the succinate semialdehyde dehydrogenase (ALDH5A1) gene and functional analysis of 27 novel disease-causing mutations in patients with SSADH deficiency. *Hum Mutat.* 2003 Dec;22(6):442-50.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/14635103>
- Blasi P, Boyl PP, Ledda M, Novelletto A, Gibson KM, Jakobs C, Hogema B, Akaboshi S, Loreni F, Malaspina P. Structure of human succinic semialdehyde dehydrogenase gene: identification of promoter region and alternatively processed isoforms. *Mol Genet Metab.* 2002 Aug;76(4):348-62.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/12208142>
- Gordon N. Succinic semialdehyde dehydrogenase deficiency (SSADH) (4-hydroxybutyric aciduria, gamma-hydroxybutyric aciduria). *Eur J Paediatr Neurol.* 2004;8(5):261-5. Review.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/15341910>

- Pearl PL, Gibson KM, Acosta MT, Vezina LG, Theodore WH, Rogawski MA, Novotny EJ, Gropman A, Conry JA, Berry GT, Tuchman M. Clinical spectrum of succinic semialdehyde dehydrogenase deficiency. *Neurology*. 2003 May 13;60(9):1413-7.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/12743223>
  - Pearl PL, Novotny EJ, Acosta MT, Jakobs C, Gibson KM. Succinic semialdehyde dehydrogenase deficiency in children and adults. *Ann Neurol*. 2003;54 Suppl 6:S73-80. Review.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/12891657>
- 

Reprinted from Genetics Home Reference:  
<https://ghr.nlm.nih.gov/gene/ALDH5A1>

Reviewed: June 2008

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications  
U.S. National Library of Medicine  
National Institutes of Health  
Department of Health & Human Services